

## Recommended Studies for Standard and Low Risk (Stages I-III) Wilms Tumor Follow-up

Time off Treatment	History and Physical Exam	Urinalysis	Creatinine SGPT, bilirubin, total protein/albumin	Dental Exam	Nuclear GFR	Chest CT scan	Chest radiograph	CT or MRI scan of abdomen and pelvis	Abdominal ultrasound	ECHO/EKG (*Reg. DD-4A Only)
End of Treatment	✓	✓	✓	✓		✓		✓		
3 months	✓	✓	✓		✓		✓		✓	
6 months	✓	✓	✓			✓		✓		
9 months	✓		✓				✓		✓	
12 months	✓	✓	✓	✓	✓	✓		✓		✓
15 months	✓						✓		✓	
18 months	✓	✓	✓			✓		✓		
21 months	✓						✓		✓	
24 months	✓	✓	✓	✓	✓	✓		✓		
27 months	✓						✓		✓	
30 months	✓	✓	✓			✓		✓		
33 months	✓						✓		✓	
36 months	✓	✓	✓	✓	✓	✓		✓		✓
42 months	✓						✓		✓	
48 months	✓	✓	✓	✓	✓		✓		✓	
54 months	✓						✓		✓	
60 months	✓	✓	✓	✓ <sup>1</sup>	✓		✓		✓	✓
Then Every 2 Years	✓	✓								✓

<sup>1</sup>Then recommended yearly throughout entire life

**If your child received abdominal radiation, the following tests are recommended:**

- **For Females:** Gynecological exam: A baseline examination at age 18 and then as indicated by your physician
- **For Males and Females:** Colonoscopy: Every five years beginning at age 35